STABLE COMPLETE MINIMAL HYPERSURFACES IN R^4 *

Haizhong Li   Guoxin Wei

Abstract

Let $x : M \rightarrow R^4$ be a stable complete minimal hypersurface, we give a sufficient condition for $x(M)$ to be a hyperplane, improving Berard's result.

1. Introduction

Let $x : M \rightarrow R^{n+1}$ be an n -dimensional minimal immersion. Let $D \subset M$ be a domain with compact closure and piecewise smooth boundary. We say that D is *stable* if it is a minimum for the volume function of the induced metric for all variations that keep the boundary D fixed. The immersion x is *stable* if any such D is stable. We will denote by A the second fundamental form of $x(M)$ in R^{n+1} , by K the scalar curvature of $x(M)$ in the induced metric.

In [2] (also see [8] or [3]), P. Berard proved the following result

Theorem 1 (see Proposition 2.2 of [2]). *Let $x : M \rightarrow R^4$ be a 3-dimensional stable complete minimal immersion such that*

$$\int_M |A|^3 dM < +\infty, \quad (1.1)$$

then $x(M) \subset R^4$ is a hyperplane.

Set

$$B_R = \{p \in M; \rho(p, p_0) \leq R\},$$

---

*This work is partially supported by the NSF of China No.10131020.

1991 Mathematics Subject Classification: 53C42, 53C40, 53A10.

Key words and phrases: stable, minimal hypersurface, hyperplane.

where ρ is the geodesic distance from a fixed point $p_0 \in M$.

In this note, we improve the above Berard's result as following:

Theorem 2. *Let $x : M \rightarrow R^4$ be a 3-dimensional stable complete minimal immersion such that*

$$\lim_{R \rightarrow \infty} \frac{\int_{B_R} |A|^3 dM}{R^{1+2q}} = 0, \quad q < \sqrt{\frac{2}{3}}, \quad (1.2)$$

then $x(M) \subset R^4$ is a hyperplane.

Remark 1. It is obvious that Berard's Theorem 1 is the direct consequence of our Theorem 2.

Remark 2. The strongly stable hypersurfaces with constant mean curvature have been studied by Alencar-do Carmo [1], do Carmo-Zhou [5] (also see survey paper of Berard-Santos [3]).

2. The proof of Theorem 2

We will follow the notation of [4] closely. Let $x : M \rightarrow R^4$ be a 3-dimensional stable complete minimal immersion.

By (2.11) of [4], we have

$$\int_M |A|^{4+2q} f^2 dM \leq \beta_1 \int_M |A|^{2+2q} |\nabla f|^2 dM, \quad (2.1)$$

where q is any positive constant with $q < \sqrt{2/3}$, β_1 is some positive constant, and f is any function on M with compact support.

We now try to transform (2.1) so that the integrand of the right hand side only involve $|A|$ in the third power. For that, we use Young's inequality ([6], p_{111}):

$$ab \leq \frac{\alpha^s a^s}{s} + \frac{\alpha^{-t} b^t}{t}, \quad \frac{1}{s} + \frac{1}{t} = 1, \quad (2.2)$$

where $\alpha > 0$ is arbitrary and $1 < s < \infty$, $1 < t < \infty$. Let p , $0 < p < 2 + 2q$, be

a number yet to be determined. By use of (2.2), we obtain

$$\begin{aligned}
 |A|^{2+2q}|\nabla f|^2 &= f^2(|A|^{2+2q}\frac{|\nabla f|^2}{f^2}) \\
 &= f^2(|A|^{2+2q-p}|A|^p\frac{|\nabla f|^2}{f^2}) \\
 &\leq f^2(\frac{\alpha^s}{s}|A|^{s(2+2q-p)} + \frac{\alpha^{-t}}{t}(|A|^p\frac{|\nabla f|^2}{f^2})^t).
 \end{aligned} \tag{2.3}$$

We now choose p to satisfy the following equations:

$$s(2+2q-p) = 4+2q, \quad pt = 3, \quad \frac{1}{s} + \frac{1}{t} = 1.$$

This is indeed possible, and the solution is

$$p = \frac{6}{1+2q}, \quad s = 1 + \frac{2}{2q-1}, \quad t = \frac{1+2q}{2}, \quad \frac{1}{2} < q < \sqrt{2/3}.$$

By use of these values and the fact that α may be made small, we obtain from (2.1) and (2.3)

$$\int_M |A|^{4+2q} f^2 dM \leq \beta_2 \int_M |A|^3 \frac{|\nabla f|^{1+2q}}{f^{2q-1}} dM, \tag{2.4}$$

where β_2 is a constant depending on q . Now we use the arbitrariness of f to replace f by $f^{\frac{1+2q}{2}}$ in (2.4) and obtain

$$\int_M |A|^{4+2q} f^{2q+1} dM \leq \beta_3 \int_M |A|^3 |\nabla f|^{2q+1} dM, \tag{2.5}$$

where β_3 is again a constant depending on q , $1/2 < q < \sqrt{2/3}$.

Choose a family of subsets B_R on M given by

$$B_R = \{p \in M : d(p, p_0) = \rho(p) \leq R\}, \tag{2.6}$$

where $\rho(p)$ is the geodesic distance on M from p to a fixed point $p_0 \in M$. By completeness, B_R is compact and $\bigcup_{R \in (0, \infty)} B_R = M$. It is well known that $|\nabla \rho| \leq 1$ almost everywhere on M (see [7]). Now fix numbers R and θ , $0 < \theta < 1$, and choose f in (2.5) by

$$\begin{aligned}
 f(p) &= 1, \quad \text{for } \rho(p) \leq \theta R, \\
 f(p) &= \frac{R - \rho(p)}{(1 - \theta)R}, \quad \text{for } \theta R \leq \rho(p) \leq R, \\
 f(p) &= 0, \quad \text{for } \rho(p) \geq R.
 \end{aligned}$$

Then from (2.5) we have

$$\int_{B_{\theta R}} |A|^{4+2q} dM \leq \beta_3 \frac{\int_{B_R} |A|^3 dM}{((1-\theta)R)^{2q+1}}, \quad \frac{1}{2} < q < \sqrt{\frac{2}{3}}.$$

By letting $R \rightarrow \infty$, we have, by assumption, that the right hand side of the above inequality approaches zero. Thus

$$\int_M |A|^{4+2q} dM = 0,$$

which implies that $|A| = 0$, i.e., $x(M) \subset R^4$ is a hyperplane. We complete the proof of Theorem 2.

3. A Related Problem

In [4], M.do Carmo and C.K.Peng proved the following well-known result

Theorem 3 (Theorem 1.3 of [4]). *Let $x : M \rightarrow R^{n+1}$ be an n -dimensional ($n \geq 3$) stable complete minimal immersion and assume that*

$$\lim_{R \rightarrow \infty} \frac{\int_{B_R} |A|^2 dM}{R^{2+2q}} = 0, \quad q < \sqrt{\frac{2}{n}}, \quad (3.1)$$

then $x(M) \subset R^{n+1}$ is a hyperplane.

By Gauss equation, we know $|A|^2 = -K$, the total curvature $\int_M |K| dM = \int_M |A|^2$, thus we have from Theorem 3

Corollary 1 (Theorem 1.2 of [4]). *Let $x : M \rightarrow R^{n+1}$ be an n -dimensional ($n \geq 3$) stable complete minimal immersion such that the total curvature is finite, that is*

$$\int_M |A|^2 dM < +\infty, \quad (3.2)$$

then $x(M) \subset R^{n+1}$ is a hyperplane.

In [8], Y. B. Shen and X. H. Zhu proved the following result, which generalized Theorem 1 for general dimension n .

Theorem 4 (Main Theorem of [8]). *Let $x : M \rightarrow R^{n+1}$ be an n -dimensional ($n \geq 3$) stable complete minimal immersion such that*

$$\int_M |A|^n dM < +\infty, \quad (3.3)$$

then $x(M) \subset R^{n+1}$ is a hyperplane.

Observing that Theorem 4 is the counterpart of Corollary 1 if we use $\int_M |A|^n dM$ instead of $\int_M |A|^2 dM$, the following problem seems to be interesting

Problem. *Can we establish the counterpart of Theorem 3 for n -dimensional ($n \geq 3$) stable complete minimal immersion $x : M \rightarrow R^{n+1}$?*

Here we would like to formulate the above problem as following:

Conjecture: *Let $x : M \rightarrow R^n$ be an n -dimensional stable complete minimal immersion and assume that*

$$\lim_{R \rightarrow \infty} \frac{\int_{B_R} |A|^n dM}{R^{1+2q}} = 0, \quad q < \sqrt{\frac{2}{n}}, \quad (3.4)$$

then $x(M) \subset R^n$ is a hyperplane.

Remark 3. Our Theorem 2 only has checked the conjecture for the case $n = 3$. In this moment, we can not get any result about the conjecture for the general case $n \geq 4$.

Acknowledgements. The first author began this research work during his visit to IME, São Paulo University from July 1 to 31, he would like to thank Professor Fabiano Brito for his hospitality. He also would like to thank Professor Paolo Piccione, chairman of the scientific committee of 13th school of differential geometry of Brazil (July 26-30 of 2004, in Sao Paulo), for inviting him to give a plenary talk in the school. Finally, the authors would like to thank referee for his or her useful comments.

References

- [1] Alencar, A., do Carmo, M. P., *Hypersurfaces with constant mean curvature in space forms*, An. Acad. Bras. Ciênc., 66(1994), 265-274.
- [2] Berard, P., *Remarques sur l'équation de J.Simons*, in Differential Geometry, A symposium in honour of Manfredo do Carmo, edited by B. Lawson and K. Tenenblat, pp. 47-57, Pitman Monograph and Surveys in Pure and Applied Mathematics 52, (1991) 47-57.
- [3] Berard, P., Santos, W., *Curvature estimates and stability properties of CMC-submanifolds in space forms*, Mat. Contemp., 17(1999), 77-97.
- [4] do Carmo, M. P., Peng, C.K., *Stable complete minimal hypersurfaces*, in Proceedings of the 1980 Beijing Symposium on Differential Geometry and Differential Equations, edited by S.S.Chern and Wu Wen-Tsun, Gordon and Breach Science Pub., Vol.3, (1982), 1349-1358.
- [5] do Carmo, M. P., Zhou, D. T., *Bernstein-type theorems in hypersurfaces with constant mean curvature*, An. Acad. Bras. Ciênc., 72(2000), 301-310.
- [6] Hardy, G. H., Littlewood, J. E., Polya, G., *Inequalities*, Cambridge University Press, (1946).
- [7] Schoen, R., Simon, L., Yau, S.-T., *Curvature estimates for minimal hypersurfaces*, Acta Math. 134(1975), 275-288.
- [8] Shen, Y. B., Zhu, X. H., *On stable complete minimal hypersurfaces in R^{n+1}* , Amer. J. Math., 120(1998), 103-116.

Department of Mathematical Sciences

Tsinghua University

Beijing, 100084

People's Republic of China

E-mail: hli@math.tsinghua.edu.cn wigx03@mails.tsinghua.edu.cn