*Matemática Contemporânea, Vol 32, 157-178*
©2007, Sociedade Brasileira de Matemática

# GLOBAL AND DECAY OF SOLUTIONS OF A DAMPED KIRCHHOFF-CARRIER EQUATION IN BANACH SPACES

**R. Izaguirre  R. Fuentes  M. Milla Miranda**

*In Homage to Professor L. A. Medeiros by his Eightieth Birthday*

**Abstract**

This paper is concerned with the study of the existence and the decay of solutions of the following problem:

$$\left| \begin{array}{l} Bu''(t) + M\left(\|u(t)\|_W^{\beta}\right) Au(t) + \delta Bu'(t) = 0, \text{ in } V', t > 0, \\ u(0) = u^0 \ , u'(0) = u^1 \ (u^0 \neq 0), \end{array} \right.$$

where A and B are symmetric linear operators from a Hilbert space V into its dual $V'$ satisfying $\langle Bv, v\rangle > 0,\ v \neq 0, \langle Av, v\rangle \geq \gamma \|v\|_V^2 , \gamma > 0$ ; W a Banach space with V continuously embedding in W; $\beta$ a real number with $\beta \geq 1$, $M(\xi)$ a smooth function with $M(\xi) \geq 0$, and $\delta$ a positive real number.

# 1 Introduction

Let V be a real separable Hilbert space whose dual is denoted by $V'$ and W a Banach space with V continuously embedding in W. Consider two symmetric linear operators $A, B : V \to V'$ such that

$$\langle Av, v\rangle \geq \gamma \|v\|_V^2 , \ \forall v \in V \ (\gamma \text{ positive constant});$$

$$\langle Bv, v\rangle > 0, \ \forall v \in V, \ v \neq 0;$$

and a smooth function $M(\xi)$ with

$$M(\xi) \geq 0, \ \forall \, \xi \geq 0.$$

*AMS Subject Classification:* 35L70,35B35
*Key Words:* Kirchhoff-Carrier Equation, global solutions, decay of solutions

Consider also two real numbers $\beta \geq 1$ and $\delta > 0$. In this conditions we have the following problem:

$$(*) \left| \begin{array}{l} Bu''(t) + M\left(\|u(t)\|_W^{\beta}\right) Au(t) + \delta Bu'(t) = 0, \text{ in } V', t > 0, \\ u(0) = u^0, \;\; u'(0) = u^1 \; (u^0 \neq 0). \end{array} \right.$$

Equation in (*) is a damped abstract version in Banach spaces of the Kirchhoff equation [14] and the Carrier equation [5]. When $B = I, \beta = 2$,W is a Hilbert space and $\delta \geq 0$, there is an extensive literature on this problem (cf. Medeiros, Limaco and Menezes [22]).

The existence of local solutions of problem (*) has been obtained by the Authors in [13].

In this paper we study the existence of global solutions of (*) when $M(\xi) \geq 0$ and the exponential decay of solutions of (*) when $M(\xi) \geq m_0 > 0$. In Section 5, we give some examples.

To obtain global solutions we use the prolongation method and in the decay of solutions, the Lyapunov approach, cf. Komornik and Zuazua [15]. In both cases it is fundamental an appropriate characterization of the derivative of $M(\|u(t)\|_W^{\beta})$. We use various results obtained in [13] and in S. S. Souza and the third A. [27].

## 2 Notations and Main Results

Let V be a real separable Hilbert space whose dual is denoted by $V'$. Consider two linear operators $A, B : V \to V'$ satisfying

$$(H1) \quad \begin{array}{l} \langle Au, v\rangle = \langle u, Av\rangle, \forall u, v \in V; \\ \langle Au, u\rangle \geq \gamma \|u\|_V^2, \forall u \in V \; (\gamma \text{ positive constant}); \\ \langle Bu, v\rangle = \langle u, Bv\rangle, \forall u, v \in V; \\ \langle Bu, u\rangle > 0, \forall u \in V, u \neq 0. \end{array}$$

Here $\langle\ ,\ \rangle$ denotes the duality pairing between $V'$ and V. We have that the scalar product $((u,v)) = \langle Au, v\rangle$ defines a norm $\|u\| = ((u,u))^{1/2}$ in V which is equivalent to the norm $\|u\|_V$. The space V will be equipped with the scalar product $((u,v))$ and norm $\|u\|$.

The bilinear form

$$(u,v) = \langle Bu, v\rangle\ ,\ \forall u, v \in V$$

is a scalar product in V. We denote by H the completed of the space $\{V,(u,v)\}$. The scalar product of the Hilbert space H will be denoted also by $(u,v)$ and its norm by $|u|$. We have that

V is densely and continuously embedding in H.

Consider the coercive self-adjoint operator S of H determined by the triplet $\{V,H,((u,v))\}$. We have:

$$(Su,v) = ((u,v)) = \langle Au, v\rangle\ ,\ \forall u \in D(S),\ \forall v \in V; \tag{2.1}$$

$$Au = BSu \text{ in } V',\ \forall u \in D(S^{3/2}). \tag{2.2}$$

Identify H with its dual $H'$. Then expression (2.1) says that A is the extension of S to the space V.

Represent by W a Banach space whose dual

(H2) $W'$ is strictly convex.

Denote by $\theta \geq 0$ a real number and by $(E_\lambda)_{\lambda\in\mathbb{R}}$ the spectral family of S. Then $D(S^\theta)$ is the Hilbert space

$$D\left(S^\theta\right) = \left\{u \in H; \int_0^\infty \lambda^{2\theta} d(E_\lambda u, u) < \infty\right\}$$

equipped with the scalar product

$$(u,v)_{D(S^\theta)} = \left(S^\theta u, S^\theta v\right).$$

Fix $\alpha \geq 0$ a real number. Assume that $D(S^{\alpha+1})$ is continuously embedding in W, that is, there exist a positive number $k_0$, such that

(H3) $$\|u\|_W \leq k_0 \|u\|_{D(S^{\alpha+1})}, \forall\, u \in D(S^{\alpha+1}).$$

Consider a function $M(\xi)$ and $\beta \geq 1$ a real number satisfying

$$(H4)\left|\begin{array}{l} M \in C^0([0,\infty[),\ M(0)=0,\ M(\xi)>0,\ \forall\,\xi>0; \\ M \in C^1(]0,\infty[); \\ |M'(\xi)|\,\lambda^{1-1/\beta} \leq C_0 M^{1/2}(\xi),\ \forall\,\xi>0\ (C_0 \text{ positive constant}). \end{array}\right.$$

Under the above considerations, we have the following result:

**Theorem 2.1** *Assume hypotheses (H1)-(H4) with $\alpha \geq 0,\ \beta \geq 1$. Consider $\delta > 0$ a real number and*

(H5) $$u^0 \in D(S^{2\alpha+5/2}), u^1 \in D(S^{2\alpha+2}), u^0 \neq 0$$

*satisfying*

(H6) $$\beta C_0 k_0 \left[\frac{\left|S^{\alpha+1}u^1\right|^2}{M\left(\|u^0\|_W^\beta\right)} + \left|S^{\alpha+3/2}u^0\right|^2\right]^{1/2} < \delta.$$

**(a)** *Then there exists a function u in the class*

$$\left|\begin{array}{l} u \in L^\infty(0,\infty; D(S^{\alpha+3/2})), \\ u' \in L^\infty(0,\infty; D(S^{\alpha+1})), \\ u'' \in L^\infty(0,\infty; D(S^{\alpha+1/2})) \end{array}\right. \tag{2.3}$$

*satisfying*

$$(P)\quad \left|\begin{array}{l} u'' + M\left(\|u\|_W^{\beta}\right) Su + \delta u' = 0 \text{ in } L^{\infty}(0,\infty; D(S^{\alpha+1/2})), \\ u(0) = u^0 \ , u'(0) = u^1. \end{array}\right.$$

**(b)** *Let $\mathcal{M}$ be the set constituted by the real numbers $T > 0$ such that there exists a unique function $u$ in the class (2.3) with $u$ solution of (P) in [0, T] and $\|u(t)\|_W > 0$ for all $t \in [0,T]$. Let $T_{max}$ be the supremum of the $T \in \mathcal{M}$. Then $\mathcal{M} \neq \emptyset$ and the solution $u$ obtained in* **(a)** *verifies*

$$u \in L^{\infty}_{loc}(0, T_{max}; D(S^{2\alpha+5/2})),$$

$$u' \in L^{\infty}_{loc}(0, T_{max}; D(S^{2\alpha+2})),$$

$$u'' \in L^{\infty}_{loc}(0, T_{max}; D(S^{\alpha+1/2}))$$

*and*

$$\frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^{\beta}\right)} + \left|S^{\alpha+3/2}u(t)\right|^2 \leq \frac{\left|S^{\alpha+1}u^1\right|^2}{M\left(\|u^0\|_W^{\beta}\right)} + \left|S^{\alpha+3/2}u^0\right|^2, \ 0 \leq t < T_{max}.$$

*And if $T_{max}$ is finite,*

$$u(t) = 0, \text{ for } t \geq T_{max}.$$

In order to obtain the decay of solutions of problem (P), we make the following considerations.

Consider a function $M(\xi)$ and $\beta > 1$ a real number satisfying

$$(H7)\quad \left|\begin{array}{l} M \in C^1([0,\infty[); \\ M(\xi) \geq m_0 > 0, \ \forall\, \xi \geq 0 \ (m_0 \text{ constant}); \\ M'(\xi) \geq 0, \ \forall\, \xi \geq 0; \\ |M'(\xi)|\, \lambda^{1-1/\beta} \leq C_1 M(\xi), \ \forall\, \xi \geq 0 \ (C_1 \text{ positive constant}). \end{array}\right.$$

As V is continuously embedding in H, we have:

$$(Su, u) = \|u\|^2 \geq C_*^2 \, |u|^2, \ \forall\, u \in D(S) \ (C_* \text{ positive constant}).$$

This implies

$$\left|S^{\alpha+1}u\right|^2 \leq \frac{1}{C_*^2}\left|S^{\alpha+3/2}u\right|^2,\ \forall\ u \in D(S^{\alpha+3/2}). \tag{2.4}$$

We introduce the constant $k_1 > 0$ verifying

$$\|u\|_W \leq k_1\left|S^{\alpha+3/2}u\right|^2,\ \forall\ u \in D(S^{\alpha+3/2}). \tag{2.5}$$

Let $\varphi(t)$ be the function

$$\varphi(t) = \frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^{\beta}\right)} + \left|S^{\alpha+3/2}u(t)\right|^2,\ t \geq 0. \tag{2.6}$$

Under the above considerations, we obtain :

**Theorem 2.2** *Assume hypotheses (H1) - (H3), (H7) with $\alpha \geq 0$ and $\beta > 1$. Consider a real number $\delta > 0$ and*

(H8) $$u^0 \in D(S^{\alpha+3/2}), u^1 \in D(S^{\alpha+1});$$

(H9) $$\beta C_1 k_0 M^{1/2}\left(k_1^{\beta}\varphi^{\beta/2}(0)\right)\varphi^{1/2}(0) < \delta;$$

*where*

$$\varphi(0) = \frac{\left|S^{\alpha+1}u^1\right|^2}{M\left(\|u^0\|_W^{\beta}\right)} + \left|S^{\alpha+3/2}u^0\right|^2.$$

*Then there exists a unique function $u$ in the class (2.3) such that $u$ is solution of Problem (P). Furthermore if*

(H10) $$\frac{\left|S^{\alpha+1}u^1\right|^2}{M\left(\|u^0\|_W^{\beta}\right)} + \left|S^{\alpha+3/2}u^0\right|^2$$

$$< min\left[\frac{\delta^2}{4\beta^2 C_1^2 k_0^2 M(k_1^{\beta}\varphi^{\beta/2}(0)}, \frac{m_0 C_*^4}{4\beta^2 C_1^2 k_0^2}, \frac{m_0 C_*^4}{4\delta^2\beta^2 C_1^2 k_0^2}\right]$$

*where $C_*$ were defined in (2.4), we have*

$$\varphi(t) \leq 3\varphi(0)e^{-\frac{\tau_0}{3}t}, \ \ t \geq 0, \tag{2.7}$$

*where*

$$\tau_0 = min(\delta, \epsilon_0), \ \epsilon_0 = min(\frac{1}{2P_0}, \frac{\delta}{4}, 1), \ P_0 = \frac{1}{C_* m_0^{1/2}} + \frac{\delta}{2C_* m_0} \tag{2.8}$$

*and $\varphi(t)$ was defined in (2.6).*

**Corollary 2.2** *If* $K = \sup_{0 \leq t < \infty} M\left(\|u(t)\|_W^\beta\right)$
*then (2.7) implies*

$$E(t) \leq 3K\varphi(0)e^{-\frac{\tau_0}{3}t}, \ \ t \geq 0,$$

*where*

$$E(t) = \left|S^{\alpha+1}u'(t)\right|^2 + M\left(\|u(t)\|_W^\beta\right)\left|S^{\alpha+3/2}u(t)\right|^2, \ t \geq 0.$$

**Remark 2.1** *By property (2.2), we have that the equations*

$$Bu''(t) + M\left(\|u(t)\|_W^\beta\right)Au(t) + \delta Bu'(t) = 0 \ in \ V', \ t > 0$$

*and*

$$u''(t) + M\left(\|u(t)\|_W^\beta\right)Su(t) + \delta u'(t) = 0 \ in \ D(S^{3/2}), \ t > 0$$

*are equivalents.*

# 3 Proof of Theorem 2.1

We need of the following result, obtained in [13]:

**Proposition 3.1** *Let $M : [0,\infty[\to \mathbb{R}$ be a function of class $C^1$ and*

$$u \in C^1([0,\infty[;W),\ u(t) \neq 0,\ \forall t \in [0,\infty[.$$

*Consider $\beta$ a real number. Then the Leibniz derivative of $M\left(\|u(t)\|_W^\beta\right)$ is given by*

$$\frac{d}{dt}\left\{M\left(\|u(t)\|_W^\beta\right)\right\} = \beta M'(\|u(t)\|_W^\beta)\|u(t)\|_W^{\beta-1}\left\langle \frac{Ju(t)}{\|u(t)\|_W}, u'(t)\right\rangle_{W'\times W},$$
$$t \geq 0,$$

*where $J$ is the duality application $J : W \to W'$ defined by*

$$\langle Jv, v\rangle_{W'\times W} = \|v\|_W^2,\ \ \|Jv\|_{W'} = \|v\|_W,\ \ \forall v \in W.$$

By [13] we have also that there exists $T_0 > 0$ and a unique function u in the class

$$\begin{aligned} &u \in L^\infty(0,T_0;D(S^{2\alpha+5/2})),\\ &u' \in L^\infty(0,T_0;D(S^{2\alpha+2})),\\ &u'' \in L^\infty(0,T_0;D(S^{2\alpha+3/2})) \end{aligned} \tag{3.1}$$

such that

$$(LP)\quad \left|\begin{array}{l} u'' + M\left(\|u\|_W^\beta\right)Su + \delta u' = 0 \text{ in } L^\infty(0,T_0;D(S^{2\alpha+3/2})),\\ u(0) = u^0,\ \ u'(0) = u^1 \end{array}\right.$$

and

$$\|u(t)\|_W > 0,\ \ \forall\, t \in [0,T_0]. \tag{3.2}$$

So $\mathcal{M} \neq \emptyset$.

Next we obtain estimates for the solutions u given in $\mathcal{M}$. Note that if u given in $\mathcal{M}$ by the uniqueness of solutions (P) in [0, T], we have that u belongs to class (3.1), u is solution of (LP) in [0, T] and u satisfies (3.2) in [0, T] (see[13]). Consider $0 < t_0 < T_{max}$. Taking the scalar product of H in both sides of equation $(LP)_1$ by $2S^{2\alpha+2}u'$, we obtain:

$$\begin{aligned} \frac{d}{dt}\left[\left|S^{\alpha+1}u'(t)\right|^2\right] &+ M\left(\|u(t)\|_W^\beta\right)\frac{d}{dt}\left[\left|S^{\alpha+3/2}u(t)\right|^2\right]\\ &+ 2\delta\left|S^{\alpha+1}u'(t)\right|^2 = 0,\ t \in [0,t_0], \end{aligned}$$

that is,

$$\frac{d}{dt}\frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)}+\frac{d}{dt}\left[\left|S^{\alpha+3/2}u(t)\right|^2\right]=-2\delta\frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)}. \tag{3.3}$$

Introduce the function

$$\varphi(t)=\frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)}+\left|S^{\alpha+3/2}u(t)\right|^2,\ \ t\in[0,t_0]. \tag{3.4}$$

Our goal is to show that $\varphi(t)$ is not increasing. By Proposition 3.1 and (3.3), we have:

$$\varphi'(t)=-\frac{1}{M\left(\|u(t)\|_W^\beta\right)}\beta M'\left(\|u(t)\|_W^\beta\right)\|u(t)\|_W^{\beta-1}$$

$$\left\langle\frac{Ju(t)}{\|u(t)\|_W},u'(t)\right\rangle_{W\times W'}\left|S^{\alpha+1}u'(t)\right|^2-\frac{2\delta\left|S^{\alpha+1)}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)},\ \ t\in[0,t_0].$$

This gives

$$\varphi'(t)\le\frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)}\left[\frac{\beta\left|M'\left(\|u(t)\|_W^\beta\right)\right|\|u(t)\|_W^{\beta-1}}{M^{1/2}(\|u(t)\|_W^\beta)}\frac{\|u'(t)\|_W}{M^{1/2}(\|u(t)|_W^\beta)}-2\delta\right].$$

Then hypothesis $(H4)_2$ and embedding (H3) give:

$$\varphi'(t)\le\frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)}\left[\beta C_0k_0\frac{\left|S^{\alpha+1)}u'(t)\right|^2}{M^{1/2}(\|u(t)|_W^\beta)}-2\delta\right],\ \ t\in[0,t_0]. \tag{3.5}$$

Introduce the function

$$\psi(t)=\beta C_0k_0\frac{\left|S^{\alpha+1)}u'(t)\right|}{M^{1/2}(\|u(t)|_W^\beta)},\ \ t\in[0,t_0].$$

We have

$$\psi(t)\le\beta C_0k_0\varphi^{1/2}(t),\ \ \forall\, t\in[0,t_0]. \tag{3.6}$$

We affirm that

$$\psi(t)<\delta,\ \ \forall\, t\in[0,t_0]. \tag{3.7}$$

In fact, suppose that there exists $t_1 \in [0, t_0]$ such that $\psi(t) \geq \delta$. By hypothesis (H6), we have $\psi(0) < \delta$. Consider

$$t^* = inf\,\{t \in [0, t_0]; \psi(t) = \delta\} > 0$$

As $\psi(t)$ is continuous in $[0, t_0]$, we have that $\psi(t^*) = \delta$, which implies by (3.5) that $\varphi(t)$ is not increasing on $[0, t^*]$. Then by hypothesis (H6) and (3.6), we obtain:

$$\psi(t) \leq \beta C_0 k_0 \varphi^{1/2}(0) < \delta, \;\; \forall\, t \in [0, t^*],$$

which is a contradiction since $\psi(t^*) = \delta$. So (3.7) holds.

If follows from (3.7), (3.5) and noting that $0 < t_0 < T_{max}$ was arbitrary that

$$\varphi'(t) \leq -\delta \frac{\left|S^{\alpha+1}u'(t)\right|^2}{M(\|u(t)|_W^\beta)}, \;\; \forall\, t \in [0, T_{max}[. \tag{3.8}$$

In particular

$$\frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)} + \left|S^{\alpha+3/2}u(t)\right|^2 \leq \frac{\left|S^{\alpha+1}u^1\right|^2}{M\left(\|u^0\|_W^\beta\right)} + \left|S^{\alpha+3/2}u^0\right|^2 <$$

$$< \frac{\delta^2}{\beta^2 C_0^2 k_0^2} = N_0^2, \;\; \forall\, t \in [0, T_{max}[. \tag{3.9}$$

Note that if $T_{max}$ is infinite then (3.9) give the theorem. Suppose that $T_{max}$ is finite. Then (3.9) implies

$$\left|S^{\alpha+3/2}u(t)\right|^2 \leq N_0^2, \;\; \forall\, t \in [0, T_{max}[. \tag{3.10}$$

As $\|u(t)\|_W \leq k_1 \|u(t)\|_{D(S^{\alpha+3/2})}$ for all $t \in [0, T_{max}[$, we have by (3.9) and (3.10) that

$$\left|S^{\alpha+1}u'(t)\right|^2 \leq N_1^2, \;\; \forall\, t \in [0, T_{max}[. \tag{3.11}$$

Consider a sequence of real number $(t_\nu)$ such that $0 < t_\nu < T_{max}$ and $t_\nu \to T_{max}$. By (3.10) and (3.11) we have that there exist $\zeta \in D(S^{\alpha+3/2})$ and

$\chi \in D(S^{\alpha+1})$ such that

$$u(t_\nu) \to \zeta \text{ weak in } D(S^{\alpha+3/2}), \tag{3.12}$$

$$u'(t_\nu) \to \chi \text{ weak in } D(S^{\alpha+1}). \tag{3.13}$$

We affirm that

$$\zeta = \chi = 0. \tag{3.14}$$

In fact, if $\zeta \neq 0$ with $\zeta$ and $\chi$ we determine the local solution w of the problem

$$\left| \begin{array}{l} w'' + M\left(\|w\|_W^\beta\right) Sw + \delta w' = 0 \text{ in } L^\infty(0, T_0; D(S^{\alpha+1/2})), \\ w(0) = \zeta \ , w'(0) = \chi \end{array} \right.$$

(see [13]). Then the function

$$\widetilde{u}(t) = \left| \begin{array}{l} w(t), \ \ 0 \leq t < T_{max} \\ w(t - T_{max}), \ \ T_{max} \leq t < T_0 + T_{max} \end{array} \right.$$

is a solution of Problem (P) in $[0, T_0 + T_{max}]$, with $\|\widetilde{u}(t)\|_W > 0$ for all $t \in [0, T_0 + T_{max}]$. This gives a contradiction with the definition of $T_{max}$. So $\zeta = 0$.

Also by (3.11),

$$\|u(t_\nu) - u(t_\mu)\|_{D(S^{\alpha+1})} \leq \int_{t_\mu}^{t_\nu} \|u'(s)\|_{D(S^{\alpha+1})} ds \leq N_1 |t_\nu - t_\mu|,$$

that is, $(u(t_\nu))$ is a Cauchy sequence in $D(S^{\alpha+1})$. As $\zeta = 0$, (3.12) implies then

$$u(t_\nu) \to 0 \ \text{ in } D(S^{\alpha+1}).$$

In particular

$$u(t_\nu) \to 0 \ \text{ in } W.$$

By estimate (3.9), we have:

$$\left|S^{\alpha+1} u'(t_\nu)\right|^2 \leq N_0^2 M\left(\|u(t_\nu)\|_W\right).$$

As $M(0) = 0$ it follows from this inequality and convergence (3.13) that $\chi = 0$. So the affirmation (3.14) is correct.

Also by equation $(P)_1$ we have that

$$u''(t_\nu) \to 0 \text{ in } D(S^{\alpha+1/2}).$$

Thus if $T_{max}$ is finite we define $u(t) = 0$ for $t \geq T_{max}$. This extension is a solution of Problem (P) in $[0, \infty[$.

□

# 4 Proof of Theorem 2.2

We begin with a previous result.

**Lemma 4.1** *Let $\beta > 1$ a real number, $M : [0, \infty[ \to \mathbb{R}$ a function of class $C^1$ and $u \in C^1([0, \infty[; W)$. Then if $u(t_0) = 0$, we have that the Leibniz derivative $\frac{d}{dt} M\left(\|u(t_0)\|_W^\beta\right)$ is equal to zero.*

**Proof:** Consider $t_0 > 0$ and $u(t_0) = 0$. Then

$$u(t_0 + h) = u(t_0 + h) - u(t_0) = h \int_0^1 u'(t_0 + \tau h) d\tau,$$

which implies for $0 < |h| < min\{1, t_0/2\}$,

$$\|u(t_0 + h)\|_W^\beta \leq |h|^\beta \left( \int_0^1 \|u'(t_0 + \tau h)\|_W d\tau \right)^\beta \leq |h|^\beta C^\beta,$$

where

$$C = max \left\{ \|u'(s)\|_W ; t_0/2 \leq s \leq t_0 + 1 \right\}.$$

The last inequality gives the result since $\beta > 1$. Analogous arguments give the result when $t_0 = 0$ and $u(0) = 0$. □

Let $u_j^0$ and $u_j^1$ be two sequences of vectors of $D(S^{2\alpha+5/2})$ and $D(S^{2\alpha+2})$, respectively, such that

$$u_j^0 \to u^0 \text{ in } D(S^{\alpha+3/2}), \quad u_j^0 \to u^1 \text{ in } D(S^{\alpha+1}).$$

By these convergence and (H9), we have:

$$\beta C_1 k_0 M^{1/2}(k_1^{\beta}\varphi_j^{\beta/2}(0))\varphi_j^{1/2}(0) < \delta, \;\; j \geq j_0,$$

where

$$\varphi_j(0) = \frac{\left|S^{\alpha+1}u_j^1\right|^2}{M(\left\|u_j^0\right\|_W^{\beta})} + \left|S^{\alpha+3/2}u_j^0\right|^2$$

Consider the problem $(j \geq j_0)$

$$(P_j) \quad \left| \begin{array}{l} u_j'' + M\left(\|u_j\|_W^{\beta}\right) Su_j + \delta u_j' = 0 \text{ in } L^{\infty}(0,\infty; D(S^{2\alpha+3/2})), \\ u_j(0) = u_j^0, \;\; u_j'(0) = u_j^1. \end{array} \right.$$

By applying the Galerkin method, the successive approximations technique and the spectral theory of the operator S, Arzela-Ascoli Theorem, Proposition (3.1), Lemma (4.1), we obtain a local solution $u_j$ of $(P_j)$, that is, we find a real number $T_0 > 0$ and a function $u_j$ in the class

$$\begin{array}{l} u_j \in \; L^{\infty}(0,T_0; D(S^{2\alpha+5/2})), \\ u_j' \in \; L^{\infty}(0,T_0; D(S^{2\alpha+2})), \\ u_j'' \in \; L^{\infty}(0,T_0; D(S^{2\alpha+3/2})) \end{array} \tag{4.1}$$

such that, $u_j$ is the solution of Problem $(P_j)$ in $L^{\infty}(0,T_0; D(S^{2\alpha+3/2}))$ (see the methodology of this approach in S. S. Souza and the third A. [27]). The same arguments allow us to obtain a real number $T_0 > 0$ and a solution u in the class

$$\begin{array}{l} u \in \; L^{\infty}(0,T_0; D(S^{2\alpha+3/2})), \\ u' \in \; L^{\infty}(0,T_0; D(S^{2\alpha+1})), \\ u'' \in \; L^{\infty}(0,T_0; D(S^{2\alpha+1/2})) \end{array}$$

such that, u is the unique solution of the problem

$$(P') \quad \left| \begin{array}{l} u'' + M\left(\|u\|_W^{\beta}\right) Su_j + \delta u' = 0 \text{ in } L^{\infty}(0,T_0; D(S^{2\alpha+1/2})), \\ u(0) = \zeta \ , u'(0) = \eta, \end{array} \right.$$

where $\zeta \in D(S^{\alpha+3/2})$ and $\eta \in D(S^{2\alpha+1})$ are arbitrary.

Let $\mathcal{M}_j$ be the set of real numbers $T > 0$ such that there exists a unique function $u_j$ in class (4.1) with T instead $T_0$ and $u_j$ is the solution of $(P_j)$ in $L^{\infty}(0,T; D(S^{2\alpha+3/2}))$. Then by the result of local existence above, we have that $\mathcal{M}_j \neq \emptyset$ . Denote by $T_{max,j}$ the supremum of $T \in \mathcal{M}$.

Let $\varphi_j(t), j \geq j_0$, be the function

$$\varphi_j(t) = \frac{\left|S^{\alpha+1}u_j'(t)\right|^2}{M\left(\|u_j(t)\|_W^{\beta}\right)} + \left|S^{\alpha+3/2}u_j(t)\right|^2, \quad t \in [0, T_{max,j}[. \tag{4.2}$$

Then as in the proof of Theorem (2.1), we obtain:

$$\varphi_j'(t) \leq \frac{\left|S^{\alpha+1}u_j'(t)\right|^2}{M\left(\|u_j(t)\|_W^{\beta}\right)} \left[\beta C_1 k_0 \left|S^{\alpha+1}u_j'(t)\right| - 2\delta\right], \quad t \in [0, T_{max,j}[. \tag{4.3}$$

Consider the function

$$\psi_1(t) = \beta C_1 k_0 \left|S^{\alpha+1}u_j'(t)\right|, \quad t \in [0, T_{max,j}[.$$

By (H9) and noting that $M(\xi)$ is increasing, we have:

$$\begin{aligned} \psi_1(0) = \quad & \beta C_1 k_0 M^{1/2}\left(\|u_j^0\|_W^{\beta}\right) \frac{\left|S^{\alpha+1}u_j^1\right|}{M^{1/2}\left(\|u_j^0\|_W^{\beta}\right)} \\ & \leq \beta C_1 k_0 M^{1/2}\left(k_1^{\beta}\varphi_j^{\beta/2}(0)\right)\varphi_j^{1/2}(0) < \delta. \end{aligned}$$

By similar arguments used in the proof Theorem 2.1, we obtain:

$$\psi_1(t) < \delta, \quad t \in [0, T_{max,j}[. \tag{4.4}$$

This result and (4.3) imply

$$\varphi_j'(t) \leq -\delta \frac{\left|S^{\alpha+1)}u_j'(t)\right|^2}{M\left(\|u_j(t)\|_W^{\beta}\right)}, \quad t \in [0, T_{max,j}[ \tag{4.5}$$

and

$$\frac{\left|S^{\alpha+1}u_j'(t)\right|^2}{M\left(\|u_j(t)\|_W^\beta\right)}+\left|S^{\alpha+3/2}u_j(t)\right|^2\leq\frac{\left|S^{\alpha+1}u_j^1\right|^2}{M\left(\|u_j^0\|_W^\beta\right)}+\left|S^{\alpha+3/2}u_j^0\right|^2<$$

$$<\frac{\delta^2}{\beta^2C_1^2k_0^2M(k_1^\beta\varphi_j^{\beta/2}(0))},\ \ t\in[0,T_{max,j}[. \tag{4.6}$$

By inequality (4.4), local existence of solution of $(P')$, uniqueness of solution of Problem $(P_j)$ in any $L^\infty(0,T;D(S^{2\alpha+3/2}))$ and by similar argument used in the proof of Theorem 2.1, we obtain that $T_{max,j}$ is infinite for $j\geq j_0$.

Fix $j\geq j_0$ and consider $\epsilon>0$. Introduce the functions

$$\rho(t)=\frac{\left(S^{\alpha+1}u_j'(t),S^{\alpha+1}u_j(t)\right)}{M\left(\|u_j(t)\|_W^\beta\right)}+\frac{\delta}{2}\frac{\left|S^{\alpha+1}u_j(t)\right|^2}{M\left(\|u_j(t)\|_W^\beta\right)},\ \ t\in[0,\infty[ \tag{4.7}$$

and

$$\varphi_\epsilon(t)=\varphi_j(t)+\epsilon\rho(t),\ \ t\in[0,\infty[, \tag{4.8}$$

where $u_j$ is the solution of $(P_j)$.

In what follows, to facilitate the notation, we omit the subscript j. We have:

$$|\rho(t)|\leq\frac{\left|S^{\alpha+1}u'(t)\right|}{C_*m_0^{1/2}M\left(\|u(t)\|_W^\beta\right)^{1/2}}\left|S^{\alpha+3/2}u(t)\right|+\frac{\delta}{2}\frac{\left|S^{\alpha+3/2}u(t)\right|^2}{m_0C_*},$$

where $C_*$ were defined in (2.4), that is,

$$|\rho(t)|\leq\left[\frac{1}{C_*m_0^{1/2}}+\frac{\delta}{2m_0C_*}\right]\varphi(t),\ \ t\in[0,\infty[.$$

So, this inequality and (4.8) imply

$$|\varphi_\epsilon(t)|\leq(1+\epsilon P_0)\,\varphi(t),\ \text{where } P_0=\frac{1}{C_*m_0^{1/2}}+\frac{\delta}{2m_0C_*}.$$

Then taking $0<\epsilon\leq 1/2P_0$, we have:

$$\frac{1}{2}\varphi(t) \leq \varphi_\epsilon(t) \leq \frac{3}{2}\varphi(t), \quad t \in [0, \infty[. \tag{4.9}$$

On the other side, by taking the scalar product of H in both sides of equation $(P_j)_1$ by $S^{2\alpha+2}u$, we obtain:

$$\left(S^{\alpha+1}u''(t), S^{\alpha+1}u(t)\right) + M\left(\|u(t)\|_W^\beta\right)\left|S^{\alpha+3/2}u(t)\right|^2 + \frac{\delta}{2}\frac{d}{dt}\left|S^{\alpha+1}u(t)\right|^2 = 0$$

or

$$\frac{\frac{d}{dt}\left(S^{\alpha+1}u'(t), S^{\alpha+1}u(t)\right)}{M\left(\|u(t)\|_W^\beta\right)} + \frac{\delta}{2}\frac{\frac{d}{dt}\left|S^{\alpha+1}u(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)} = \frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)} - \left|S^{\alpha+3/2}u(t)\right|^2$$

Combining this equality with the definition (4.7) of $\rho(t)$, we deduce, for $u(t) \neq 0$,

$$\rho'(t) = \frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)} - \left|S^{\alpha+3/2}u(t)\right|^2 -$$

$$\frac{\beta M'\left(\|u(t)\|_W^\beta\right)\|u(t)\|_W^{\beta-1}}{M\left(\|u(t)\|_W^\beta\right)}\left\langle\frac{Ju(t)}{\|u(t)\|_W}, u'(t)\right\rangle$$

$$\left(\frac{S^{\alpha+1}u'(t)}{M^{1/2}\left(\|u(t)\|_W^\beta\right)}, \frac{S^{\alpha+1}u(t)}{M^{1/2}\left(\|u(t)\|_W^\beta\right)}\right) -$$

$$\frac{\delta\beta M'\left(\|u(t)\|_W^\beta\right)\|u(t)\|_W^{\beta-1}}{2M\left(\|u(t)\|_W^\beta\right)}\left\langle\frac{Ju(t)}{\|u(t)\|_W}, u'(t)\right\rangle\frac{\left|S^{\alpha+1}u(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)} =$$

$$= \frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)} - \left|S^{\alpha+3/2}u(t)\right|^2 - L_1 - L_2; \tag{4.10}$$

and for $u(t) = 0$,

$$\rho'(t) = \frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)} - \left|S^{\alpha+3/2}u(t)\right|^2 \tag{4.11}$$

(see Lemma 4.1). We have, by hypothesis $(H7)_4$, (4.4), and hypothesis $(H10)$:

$$|L_1| \le \beta C_1 k_0 \left|S^{\alpha+1}u'(t)\right| \frac{\left|S^{\alpha+1}u'(t)\right|}{M^{1/2}\left(\|u(t)\|_W^\beta\right)} \frac{\left|S^{\alpha+1}u(t)\right|}{M^{1/2}\left(\|u(t)\|_W^\beta\right)} \le$$

$$\le \beta C_1 k_0 \left|S^{\alpha+1}u'(t)\right| \left(\frac{\left|S^{\alpha+1}u'(t)\right|^2}{2M\left(\|u(t)\|_W^\beta\right)} + \frac{\left|S^{\alpha+1}u(t)\right|^2}{2M\left(\|u(t)\|_W^\beta\right)}\right) \le$$

$$\le \frac{\delta}{4}\frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)} + \frac{\beta C_1 k_0 \left|S^{\alpha+1}u'(t)\right| \left|S^{\alpha+3/2}u(t)\right|^2}{2m_0^{1/2}C_*^2 M\left(\|u(t)\|_W^\beta\right)^{1/2}} \le$$

$$\le \frac{\delta}{4}\frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)} + \frac{1}{4}\left|S^{\alpha+3/2}u(t)\right|^2.$$

Also

$$|L_2| \le \frac{\delta C_1 k_0}{2m_0^{1/2}C_*^2}\frac{\left|S^{\alpha+1}u'(t)\right|}{M^{1/2}\left(\|u(t)\|_W^\beta\right)}\left|S^{\alpha+3/2}u(t)\right|^2 \le \frac{1}{4}\left|S^{\alpha+3/2}u(t)\right|^2.$$

Combining (4.10), (4.11) and the last two inequalities, we obtain:

$$\rho'(t) \le \frac{\left|S^{\alpha+1}u'(t)\right|}{M\left(\|u(t)\|_W^\beta\right)} - \left|S^{\alpha+3/2}u(t)\right|^2 + \frac{\delta}{4}\frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)} + \frac{1}{2}\left|S^{\alpha+3/2}u(t)\right|^2.$$

This inequality, the definition (4.8) of $\varphi_\epsilon$ and inequality (4.3), give:

$$\begin{aligned}\varphi_\epsilon'(t) \quad &\le -\delta\frac{\left|S^{\alpha+1}u'(t)\right|}{M\left(\|u(t)\|_W^\beta\right)} + \epsilon\frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)} - \epsilon\left|S^{\alpha+3/2}u(t)\right|^2 \\ &+\epsilon\frac{\delta}{4}\frac{\left|S^{\alpha+1}u'(t)\right|^2}{M\left(\|u(t)\|_W^\beta\right)} + \quad \frac{\epsilon}{2}\left|S^{\alpha+3/2}u(t)\right|^2,\end{aligned}$$

for all $u(t), t \in [0, \infty[$. Noting that $\epsilon \leq min\{1, \delta/4\}$, we obtain:

$$\varphi_\epsilon'(t) \leq -\frac{\delta}{2} \frac{\left|S^{\alpha+1}u'(t)\right|}{M\left(\|u(t)\|_W^\beta\right)} - \frac{\epsilon}{2}\left|S^{\alpha+3/2}u(t)\right|^2, \ \ t \in [0, \infty[,$$

that implies by (4.9),

$$\varphi_\epsilon'(t) \leq -\frac{\tau_0}{3}\varphi_\epsilon(t), \ \ t \in [0, \infty[,$$

which gives

$$\varphi_\epsilon(t) \leq \varphi_\epsilon(0)e^{-\frac{\tau_0}{3}t}, \ \ t \in [0, \infty[.$$

Therefore, by (4.9),

$$\varphi_j(t) \leq 3\varphi_j(0)e^{-\frac{\tau_0}{3}t}, \ \ t \in [0, \infty[, \ \ j \geq j_0.$$

By the methodology used in [27], we obtain the limit u of the solutions $u_j$ is the solution of Problem (P) with u in class (2.3). Also by taking the lim inf in both sides of the last inequality, we deduce inequality (2.7).

□

# 5 Examples

The result obtained in Theorem 2.1 can be applied to the equation

$$u''(t) + \|u(t)\|_W^2 Su(t) + \delta u'(t) = 0, \ \ t > 0.$$

Here $M(\xi) = \xi^{2/\beta}$. And the result obtained in Theorem 2.2, to the equation

$$u''(t) + M\left(\|u(t)\|_W^\beta\right) Su(t) + \delta u'(t) = 0, \ \ t > 0,$$

where $M(\xi) = \xi^\sigma + m_0$, $\sigma \geq 1/\beta$ and $m_0 > 0$. For the mixed problem associated to these equation, see [13].

**REFERENCES**

[1] Arosio, A.; Garavaldi, S., *On the mildly Kirchhoff string*, Math. Appl. Science, 14 (1991), 177-195.

[2] Arosio, A.; Spagnolo, S., *Global solutions to the Cauchy problem for a nonlinear hyperbolic equation*, Nonlinear Partial Differential Equations, H. Brezis and J. l.Lions eds., College de France Seminar, Vol. 6, Pitman, London, (1984), 1-26.

[3] Bernstein, S., *Sur une classe d'équations functionelles aux derivées partielles*, Izv. Acad. Nauk SSSR, Ser. Math. , 4 (1940), 17-26.

[4] Brezis, H., *Analyse Fonctionelle, Théorie et Applications*, Masson, Paris, (1983).

[5] Carrier, G. F., *On the non-linear vibration problem of the elastic string*, Quart. Appl. Math., 3 (1945), 157-165.

[6] Clark, H. R., *Global classical solutions to the Cauchy problem for a nonlinear wave equation*, Internat. J. Math. & Math. Sci., 21(1998), 533-548.

[7] Corrêa, F. J. S. A.; Nascimento, R. G., *On an elliptic equation with a Kirchhoff term*, Proceedings 62 SBA, UNIRIO, Rio, (2005), 1-8.

[8] Cousin, A. T.; Frota, C. L.; Larkin, N. A.; Medeiros, L. A., *On the abstract model of Kirchhoff-Carrier equation*, Computational and Appl. Analyisi, 1 (1997), 389-404.

[9] Crippa, H. R., *On local solutions of some mildly degenerate hyperbolic equation*, Nonlinear Analysis : TMA, 21 (1993), 565-574.

[10] Ebihara, Y.; Medeiros, L. A.; Milla Miranda, M., *Local solutions for a nonlinear degenerate hyperbolic equation*, Nonlinear analysis :TMA, 10 (1986), 27-40.

[11] Ekeland, I.; Teman, R., *Convex Analysis and Variational Problems*, North-Holland, Amsterdam, (1976).

[12] Frota, C. L., *Non local solutions of a nonlinear hyperbolic partial differential equation*, Portugaliae Mathematica, 51 (1994), 455-473.

[13] Izaguirre, R.; Fuentes, R.; Milla Miranda, M., *Existence of local solutions of the Kirchhoff-Carrier equation in Banach spaces*, to appear in Nonlinear Analysis: TMA.

[14] Kirchhoff, G., *Vorlesunger über Mechanik*, Teubner, Leipzig, (1883).

[15] Kormonik, V.; Zuazua, E., *A direct method for boundary stabilization of the wave equation*, J. Math. Pure et Appl., 69 (1990), 33-54.

[16] Larkin, N. A., *Global regular solution for the nonhomogeneous Carrier equation*, Mathematical Problems in Enginering, 8 (2002), 15-31.

[17] Limaco, J.; Clark, H. R.; Medeiros, L. A., *On damped Kirchhoff equation with variable coefficients*, J. Math. Anal. Appl., 307 (2005), 641-655.

[18] Lions, J.-L., *Quelques Méthodes de Résolution de Problèmes aux Limites Non Linéaires*, Dunod, Paris, (1969).

[19] Lions, J.-L., *On some equations in boundary value problems of mathematical physics*, Contemporary Developmemt in Continuous Mechanics and Partial Differential Equations, G. de la Penha and L. A. Medeiros eds., North-Holland, London, (1978).

[20] Matos, M. P., *Mathematical analysis of the nonlinear model for the vibrations of a string*, Nonlinear Analysis : TMA, 17 (1991), 1125-1137.

[21] Medeiros, L. A.; Milla Miranda, M., *Solutions for the equation of nonlinear vibrations in Sobolev spaces of fractionary order*, Mat. Aplic. Comp., 6 (1987), 257-276.

[22] Medeiros, L.A.; Limaco, J.; Menezes, S. B., *Mathematical vibrations of elastic string : mathematical aspects*, part one, J. Comp. Analysis and Applications, 4 (2002), 91-127.

[23] Nishihara, K., *On a global solutions of some quasilinear hyperbolic equation*, Tokyo, J. Math., 7(1984), 437-459.

[24] Pohozaev, S. I., *On a class of quasilinear hyperbolic equations*, Math. Sbornik, 95 (1975), 152-166.

[25] Pohozaev, S. I., *The Kirchhoff quasilinear hyperbolic equation*, Differential Equations, 21 (1985), 101-108.

[26] Rivera, P. H., *On local strong solution of a nonlinear partial differential equation*, Applicable Analysis, 10 (1980), 93-104.

[27] Souza, S. S.; Milla Miranda, M., *Existence and decay of solutions of a damped Kirchhoff equation*, Int. J. of Pure and Appl. Math. 32 (2006), 483-508.

[28] Yamada, Y., *Some nonlinear degenerate wave equation*, NonlinearAnalysis : TMA, 11 (1987), 1155-1168.

[29] Yamazaki, T., *On local solutions of some quasilinear degenerate hyperbolic equations*, Funkciakj Ervacioj, 31 (1988), 439-457.

[30] Zuazua, E., *Controlabilidad Exacta yEstabilización dela Ecuación de Ondas*, Textos de Métodos Matemáticos, 23, IM-UFRJ, Rio, (1991).

[31] Zeidler, E., *Nonlinear Functional Analysis and Its Applications*, Vol. III, Springer-Verlag, New York, (1985).

Facultad de Ciencias Matemáticas
UNMSM
Av. Venezuela S/N,
Lima, PERU

Instituto de Matemática - UFF
Rua Mário Santos Braga S/N
Valonguinho
24.020-140, Niterói, RJ, BRAZIL
*E-mail*: ricardof16@yahoo.com.br

Instituto de Matemática - UFRJ
Caixa Postal 68530
21945-970, Rio de Janeiro, RJ, BRAZIL
*E-mail*: milla@im.ufrj.br